

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

AM

31/8



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/445, 9/14, 9/16	A1	(11) International Publication Number: WO 98/31365 (43) International Publication Date: 23 July 1998 (23.07.98)
(21) International Application Number: PCT/GB98/00081 (22) International Filing Date: 12 January 1998 (12.01.98) (30) Priority Data: 9700692.8 15 January 1997 (15.01.97) GB 9714873.8 15 July 1997 (15.07.97) GB (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): JACEWICZ, Victor, Witold [GB/GB]; SmithKline Beecham Pharmaceuticals, Old Powder Mills, Near Leigh, Tonbridge, Kent TN11 9AN (GB). WARD, Neal [GB/GB]; SmithKline Beecham Pharmaceuticals, Old Powder Mills, Near Leigh, Tonbridge, Kent TN11 9AN (GB). (74) Agent: GIDDINGS, Peter, John; SmithKline Beecham plc, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PAROXETINE COMPOSITIONS		
(57) Abstract Paroxetine hydrochloride is obtained in a free-flowing and easily soluble form (suitable for preparing solid formulations or aqueous solutions, suitable for parenteral use) by spray-drying solutions of paroxetine hydrochloride hemihydrate or other anhydrate hydrate solvate amorphous forms.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PAROXETINE COMPOSITIONS

The present invention relates to a process for the preparation of a pharmaceutically active compound, and to use of the so-prepared compound in therapy. In particular this invention
5 is concerned with the preparation of a free-flowing form of paroxetine hydrochloride.

Pharmaceutical products with antidepressant and anti-Parkinson properties are described in US-A-3912743 and US-A-4007196. An especially important compound among those disclosed is paroxetine, the (-)-*trans* isomer of 4-(4'-fluorophenyl)-3',4'-methylenedioxy-
10 phoxymethyl)-piperidine. This compound is used in therapy as the hydrochloride salt to treat *inter alia* depression, obsessive compulsive disorder (OCD) and panic.

Paroxetine hydrochloride has been described in the literature as a crystalline hemihydrate (see EP-A-0223403 of Beecham Group) and as various crystalline anhydrate forms (see
15 WO96/24595 of SmithKline Beecham plc). These known forms have properties that are not ideal for all pharmaceutical applications, and are prepared by multi-step procedures involving precipitation under carefully controlled conditions, filtration, drying, and homogenisation. The preferred crystallisation procedures utilise organic solvents which, when compared to water, are costly and are associated with safety and environmental
20 problems. Furthermore, the difficulty of producing crystalline products with a uniform and regular particle size causes problems with formulation by encapsulation. Also, the flow characteristics of crystalline products limit the choice of bulk transfer and formulation technologies that can be used, while dust formation and electrostatic properties can be hazardous. In addition, the known sold forms of paroxetine hydrochloride are relatively
25 insoluble and are slow to dissolve completely.

There remains a need for a form of paroxetine hydrochloride with improved processing and formulation characteristics.

30 According to a first aspect of the invention, there is provided a process for preparing a free-flowing form of paroxetine hydrochloride which comprises spray drying a solution of paroxetine hydrochloride.

The feedstock for spray drying may be prepared conveniently by, for example, dissolution
35 of paroxetine free base in aqueous hydrochloric acid, although other solid forms of
paroxetine hydrochloride may also be dissolved. For example, the feedstock may be
prepared by dissolving amorphous paroxetine hydrochloride or a crystalline paroxetine
hydrochloride anhydrate, hydrate or solvate in suitable solvent. The solvent used may be

pure water or a mixture of water with compatible organic solvents. Suitable compatible organic solvents include pyridinem acetic acid, acetonitrile, acetone, ethanol, propan-1-ol, butan-1-ol and tetrahydrofuran. Or alternatively a suitable organic solvent may be used on its own to form a solution with paroxetine hydrochloride. Some heating may be used to achieve and maintain complete solution, though once dissolved and in the absence of seeds of a crystalline form, aqueous solutions are stable at ambient temperature for many days. Suitable concentrations of paroxetine hydrochloride for spray-drying are in the range 1 to 30% by weight, preferably in the range 5% to 20% by weight.

- 10 Using conventional spray-drying procedures under normal conditions, often results in paroxetine hydrochloride particles that are sticky and adhere to the sides of the apparatus and to each other. However, when apparatus and operating conditions are selected to ensure that the particles are cooled sufficiently before they strike the apparatus walls, successful spray-drying may be carried out. Careful control of drop size in the spray
15 nozzles, air flow rates and temperatures is needed to suit the apparatus used.

The paroxetine product of the above process is free-flowing, is readily wetted, and dissolves rapidly; solutions with high concentrations may be prepared without recourse to heating.

20

Accordingly, a second aspect of this invention is spray-dried paroxetine hydrochloride.

- Spray-dried paroxetine hydrochloride of this invention has been found to be particularly suitable for applications where uniform particle size and good flow properties are
25 advantageous. Furthermore as a result of the close control of particle size possible by spray-drying, the product may be handled conveniently and safely without the hazards associated with the dust produced when conventionally prepared paroxetine hydrochloride solids are prepared. Examples of applications where uniform particle size are advantageous include controlled release and microencapsulation (coated particle
30 technology). Samples may be produced with particle sizes for specific applications, for example in the range 10-1000 microns.

- Microencapsulation may be incorporated into the spray-drying process or may be carried out in a subsequent step. This technology is useful for taste masking, rapid or controlled
35 release formulations, hence control of pharmacokinetics including the matching of pharmacokinetic properties for combination products.

Isolation of the solid product from the feedstock solution may be possible with just one processing stage; and so there is generally no need for blending, granulating, or drying, though an extra drying stage may be added if required. Providing aqueous feedstocks are used the costs and environmental problems normally associated with organic solvents are entirely avoided.

The spray-dried product of this invention may be formulated for therapy in the dosage forms described in EP-A-0223403 or WO96/24595. The free-flowing properties are advantageous for the preparation of solid formulations. Also the easily soluble nature of spray dried paroxetine hydrochloride makes it suitable for the preparation of solutions for parenteral use.

Therapeutic uses of the paroxetine product of this invention include treatment of: alcoholism, anxiety, depression, obsessive compulsive disorder, panic disorder, chronic pain, obesity, senile dementia, migraine, bulimia, anorexia, social phobia, pre-menstrual syndrome (PMS), adolescent depression, trichotillomania, dysthymia, and substance abuse, referred to below as "the disorders".

Accordingly, the present invention also provides:

a pharmaceutical composition for treatment or prophylaxis of the disorders comprising spray-dried paroxetine hydrochloride and a pharmaceutically acceptable carrier or an aqueous solution of reconstituted spray-dried paroxetine hydrochloride;

the use of spray-dried paroxetine hydrochloride to manufacture a medicament in solid or reconstituted liquid form for the treatment or prophylaxis of the disorders; and

a method of treating the disorders which comprises administering an effective or prophylactic amount of spray-dried paroxetine hydrochloride as a solid oral composition or as a reconstituted aqueous oral or parenteral composition to a person suffering from one or more of the disorders.

The invention is illustrated by the following Example..

Example:

- 5 A 10% aqueous solution of paroxetine hydrochloride is spray-dried under the following conditions:

	Apparatus:	Niro Fielder Mobile Minor
	Inlet temperature setting:	185°C
10	Actual inlet temperature:	184-185°C
	Outlet temperature:	94-95°C
	Atomiser speed:	40,000 - 50,000 rpm
	Pump speed (peristaltic):	32-34 rpm
	Air supply	4.8 - 5.2 bar
15	DP across filters:	
	Bag filter:	start of run 57 mm of water end of run 65 mm of water
	Hepa filter:	start of run 7 mm of water end of run 7 mm of water
20	DP across the orifice plate:	start of run 80+ mm of water end of run 80+ mm of water

CLAIMS

1. A process for preparing a free-flowing form of paroxetine hydrochloride which comprises spray drying a solution of paroxetine hydrochloride.
- 5 2. A process according to claim 1, in which the feedstock for spray drying is prepared by dissolution of paroxetine free base in aqueous hydrochloric acid.
- 10 3. A process according to claim 1, in which the feedstock is prepared by dissolving amorphous paroxetine hydrochloride or a crystalline paroxetine hydrochloride anhydrate, hydrate or solvate in a suitable solvent.
- 15 4. A process according to claim 1,2 or 3, in which the solvent is pure water or a mixture of water with one or more compatible organic solvents.
- 20 5. A process according to claim 1 or 3 in which the solution of paroxetine hydrochloride is in a suitable organic solvent in the absence of water.
6. A process according to claim 4 or 5 in which the organic solvent is selected from pyridine, acetic acid, acetonitrile, acetone, ethanol, propan-1-ol, butan-1-ol, or tetrahydrofuran
- 25 7. A process according to any one of the preceding claims, wherein the concentration of paroxetine hydrochloride is in the range 5% to 20% by weight.
- 30 8. Spray-dried paroxetine hydrochloride.
9. A pharmaceutical composition for treatment or prophylaxis of the disorders comprising spray-dried paroxetine hydrochloride and a pharmaceutically acceptable carrier or an aqueous solution of reconstituted spray-dried paroxetine hydrochloride.
10. The use of spray-dried paroxetine hydrochloride to manufacture a medicament in solid or reconstituted liquid form for the treatment or prophylaxis of the disorders.
- 35 11. A method of treating the disorders which comprises administering an effective or prophylactic amount of spray-dried paroxetine hydrochloride as a solid oral composition or as a reconstituted aqueous oral or parenteral composition to a person suffering from one or more of the disorders.

12. A composition according to claim 9, use according to claim 10, or a method according to claim 11, wherein the spray-dried paroxetine hydrochloride is the product of a process claimed in any one of claims 1 to 7.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 98/00081

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/445 A61K9/14 A61K9/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	EP 0 810 224 A (ASAHI) 3 December 1997 see claims see examples	1,2,5,6, 8
A	GB 2 297 550 A (SMITHKLINE BEECHAM) 7 August 1996 cited in the application see the whole document	1-12

☐ Further documents are listed in the continuation of box C

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

18 May 1998

Date of mailing of the international search report

27.05.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Scarponi, U

INTERNATIONAL SEARCH REPORT

...formation on patent family members

International Application No

PCT/GB 98/00081

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 810224	A	03-12-1997	CA 2206592 A	30-11-1997
			JP 10045756 A	17-02-1998
<hr/>				
GB 2297550	A	07-08-1996	CY 2015 A	20-02-1998
			AU 4332896 A	15-08-1996
			AU 4786496 A	27-08-1996
			BE 1009112 A	05-11-1996
			BG 100333 A	30-08-1996
			BR 9600534 A	13-05-1997
			CA 2168829 A,C	07-08-1996
			CA 2210022 A	07-08-1996
			CA 2211521 A	07-08-1996
			CA 2211522 A	07-08-1996
			CH 688353 A	15-08-1997
			CN 1143643 A	26-02-1997
			CZ 9600320 A	14-08-1996
			DE 19603797 A	14-08-1996
			DK 11996 A	07-08-1996
			WO 9624595 A	15-08-1996
			EP 0808314 A	26-11-1997
			FI 960519 A	07-08-1996
			FR 2730232 A	09-08-1996
			GR 1002466 B	06-11-1996
			HK 59397 A	16-05-1997
			HU 9600255 A	28-03-1997
			IE 960104 A	07-08-1996
			IT MI960203 A	05-08-1997
			JP 8245620 A	24-09-1996
			LT 96007 A,B	25-10-1996
			LU 88711 A	23-08-1996
			LV 11618 B	20-04-1997
			MC 2411 A	02-12-1996
			NL 1002248 C	11-09-1996
			NL 1002248 A	06-08-1996
			NO 960472 A	07-08-1996
			NZ 280943 A	29-01-1997
			PL 312646 A	19-08-1996
			PT 101827 A,B	30-09-1996
			SE 9600406 A	07-08-1996
			SG 43787 A	14-11-1997

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/00081

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 2297550 A		SI 9600036 A	31-10-1996
		SK 14396 A	06-11-1996
		NO 970939 A	07-08-1996
